

Table of Contents

Preface	vⁱⁱ
Organization	ix
Opening Address	xil
<i>H. Sakai</i>	
Welcome Greetings	xiⁱⁱ
<i>H. Suematsu</i>	
Welcome Speech	x^v
<i>W. Iwamoto</i>	
GTGR	
History of giant resonances and quenching	3
<i>A. Arima</i>	
Toward precise Gamow-Teller strengths from charge exchange data	14
<i>C. D. Goodman</i>	
Quenching problem in Gamow-Teller giant resonance	29
<i>H. Sakai, T. Wakasa, H. Okamura, T. Nonaka, T. Ohnishi, K. Yako, K. Sekiguchi, S. Fujita, Y. Satou, H. Otsu, T. Uesaka, S. Ishida, N. Sakamoto, M. B. Greenfield and K. Hatanaka</i>	
Spin polarization transfer in the (\vec{p}, \vec{n}) and $(\vec{\text{He}}, \vec{t})$ reactions	4 0
<i>B. T. Kim, H. S. Lee and T. Udagawa</i>	
Gamow-Teller strengths in $(d, {}^2\text{He})$ reactions	5 0
<i>C. A. Gagliardi, G. K. Ajupova, A. C. Betker, B. C. Hyman, Y.-W. Lui, J. R. Musser, R. E. Tribble and H. M. Xu</i>	
Gamow-Teller states observed in the $(d, {}^2\text{He})$ reactions on intermediate-mass nuclei	6 0
<i>T. Niizeki, K. Kato, T. Yamamoto, T. Yamashita, T. Ichihara, S. Ishida, H. Otsu, N. Sakamoto, T. Uesaka, T. Wakasa, K. Itoh, H. Orihara, Y. Tajima, H. Ohnuma, H. Okamura, H. Sakai and Y. Satou</i>	
The (n, p) reaction — not so boring after all?	7 0
<i>J. Blomgren</i>	

Gamow-Teller strengths from charge-exchange reactions on light nuclei and significance of alpha cluster	80
<u>M. Fjiwara and I. Daito</u>	
QES and RL/RT	
Spin response functions studied by quasi-elastic scattering	93
<u>M. Ichimum, K. Nishida, A. Itabashi and K. Kawahigashi</u>	
Polarization transfer in quasifree (\vec{p}, \vec{n}) reactions at 200 MeV	106
J. Rapaport	
Nuclear spin responses from polarization transfer in quasi-free (p, n) reactions at 346 MeV	117
<u>T. Wakasa, H. Otsu, S. Ishida, N. Sakamoto, T. Uesaka, H. Sakai, H. Okamura, T. Nonaka, Y. Satou, G. Yokoyama, S. Fujita, M. B. Greenfield, K. Hatanaka and M. Ichimum</u>	
Spin-isospin excitations in exotic nuclei	
Gamow-Teller P-decays in drip line nuclei	129
<u>I. Hamamoto</u>	
Prospecting along the N = Z line via delayed proton emission	140
<u>C. Micke, J. Giovinazzo, Ph. Dessagne, A. Huck, A. Knipper, G. Marguier, C. Longour, V. Rauch, G. Walter, M. J. G. Borge, O. Tengblad, A. Jokinen and M. Ramdhane</u>	
The GT resonance revealed in β^+ -decay using new experimental techniques	150
<u>J. Agmmunt, A. Algom, D. Cano-Ott, A. Gadea, B. Rubio, J. L. Tain, M. Gierlik, M. Karny, Z. Janas, A. Piochocki, K. Rykaczewski, J. Szerypo, R. Collatz, J. Gerl, M. Górska, H. Grawe, M. Hellström, Z. Hu, R. Kirchner, M. Rejmund, E. Roeckl, M. Shibata, L. Batist, F. Moroz, V. Wittmann and P. Kleinheinz</u>	
Spin-isospin degrees of freedom in unstable nuclei	160
<u>T. Otsuka and T. Suzuki</u>	
Spin dipole and higher spin-multipole GR	
Spin-dipole resonance studied by the $(\vec{d}, {}^2\text{He})$ reaction	173
<u>H. Okamura, T. Wakasa, Y. Satou, T. Ohnishi, T. Nonaka, H. Sakai, S. Ishida, N. Sakamoto, T. Uesaka, T. Ichihara, T. Niizeki and H. Ohnuma</u>	

Spin dipole states of deformed nuclei	183
H. Kurasawa and T. Suzuki	
Spin-dipole states in ^{12}C and ^{16}O	191
H. Sagawa and T. Suzuki	
GT strength in odd-A nuclei	203
J. W. Watson	
Recent results on the magnetic dipole and quadrupole response in nuclei probed in electron and photon scattering	213
A. Richter	
Isovector resonances in light nuclei studied by the ($^7\text{Li}, ^7\text{Be}$) reaction	238
S. Nakayama, K. Fushimi, N. Koori, H. Akimune, Y. Arimoto, I. Daito, H. Fujimura, M. Fujiwara, T. Inomata, K. Ishibashi, K. Takahisa, H. Yoshida, M. Tanaka, Y. Fujita, H. Kohri, T. Takeuchi, T. Yamagata, Y. Yamamoto, K. Yoneham and A. Tamii	
Microscopic structure of SGR	
What can we learn from particle- and gamma-decay of charge-exchange giant resonances?	247
M. N. Hamkeh, H. Akimune, Y. Arimoto, I. Daito, H. Ejiri, H. Fujimura, Y. Fujita, M. Fujiwam, K. Fushimi, N. Ihara, T. Inomata, K. Ishibashi, J. Jiinecke, H. Kohri, N. Matsuoka, H. K. T. Van Der Molen, S. Nakayama, N. Takamum, A. Tamii, M. Tanaka, H. Toyokawa, M. Yoshimura and M. Yosoi	
Isospin mixing in the isobaric-analog state derived from the ($p, n_{IAS}\bar{p}$) reaction on ^{140}Ce , $^{172,174,176}\text{Yb}$ and ^{208}Pb	264
H. Oriham, C. C. Yun, A. Temkawa, K. Itoh, A. Yamamoto, K. Kawami, H. Suzuki, Y. Mizuno, H. Kamurai, K. Ishii, Y. Fujii, H. Sagawa, T. Suzuki and H. Ohnuma	
Proton decays from spin-isospin excitations via the ($^3\text{He} t$) reactions at 450 MeV 	274
H. Akimune, I. Daito, H. Fujimura, Y. Fujita, M. Fujiwara, M. N. Hamkeh, F. Ihara, J. Jiinecke, K. Ishibashi, H. Kohri, S. Nakayama, A. Tamii, M. Tanaka, H. Toyokawa and M. Yosoi	
Widths of Gamow-Teller and isobaric analog resonances: microscopic approach	280
G. Colo, P. F. Bortignon, H. Sagawa, N. Van Gai and T. Suzuki	

Isobaric analog state of "Li (experiment)	295
<u>T. Temnishi, S. Shimoum, Y. Ando, M. Hirai, N. Iwasa,</u>	
<u>T. Kikuchi, S. Moriya, T. Motobayashi, H. Mumkami, T. Nakamum,</u>	
T. Nishio, <u>H. Sakumi, T. Uchibori</u> , Y. Watanabe, Y. Yanagisawa and <u>M. Ishiham</u>	
Isobaric analog state of ^{11}Li	302
<u>T. Suzuki and T. Otsuka</u>	
Structure and decay properties of SDR within semimicroscopical approach	312
<u>E. A. Moukhai, V. A. Rodin</u> and <u>M. H. Urin</u>	
Fine structure and isospin structure of GT and MI resonances	322
<u>Y. Fujita</u>	
Spin-isospin excitations in related Aeld	
Nuclear spin responses for neutrinos and nuclear spin giant resonances ..	335
<u>H. Ejiri</u>	
Spin strength and supernovae	347
<u>S. M. Austin</u>	
Deeply bound pionic atoms — new nuclear resonances of Gamow-Teller type	357
<u>T. Yamazaki</u>	
Effective interactions	
Medium effects on the NN interaction from (p, p) reactions: evidence for chiral restoration?	369
<u>E. J. Stephenson and F. Sammarruca</u>	
NN interactions in nuclear field studied by $(p, 2p)$ reactions	378
<u>T. Noro, H. Akimune, T. Baba, I. Daito, K. Hatanaka,</u>	
<u>M. Kawabata, F. Ihara, Y. Maeda, N. Matsuoka</u> , Y. Mizuno,	
K. Takahisa, <u>K. Tamum, M. Tanaka, H. Toyokawa, H. Yoshida,</u>	
<u>M. Yoshimum, Y. Yuasa, M. Ito</u> , <u>T. Kawabata, M. Nakamum,</u>	
<u>H. Sakaguchi, H. Takeda, T. Taki</u> , A. Tamii, <u>S. Toyama, M. Yosoi,</u>	
<u>H. Akiyoshi, S. Morinobu, K. Sagara, A. Okihana</u> and <u>H. Yamazaki</u>	
Isoscalar spin excitations	
RPA/DWIA description of spin excitations in the continuum by (\vec{p}, \vec{p}') and (\vec{d}, \vec{d}') reactions	391
<u>F. T. Baker</u>	

One- and two-phonon spin-dependent excitations in polarized deuteron scattering	403
T. Suzuki	
Study of isoscalar spin strength via the $^{12}\text{C}(d, d')$ reaction	414
S. Ishida, N. Sakamoto, T. Ichiham, Y. Satou, H. Sakai, H. Okamura, H. Otsu, T. Uesaka, T. Wakasa, T. Ohnishi, T. Nonaka, G. Yokoyama, T. Niizeki and N. Nishimori	
Spin excitations by $^{12}\text{C}(\vec{p}, \vec{p}')$ reactions at $\theta = 0$ degrees	424
A. Tamii, T. Ishikawa, M. Itoh, T. Kawabata, M. Nakamura, H. Sakaguchi, H. Takeda, T. Taki, M. Yosoi, H. Akimune, I. Daito, M. Fujiwara, K. Hatanaka, F. Ihara, K. Ishibashi, T. Inomata, T. Noro, E. Obayashi, H. Yoshida, M. Yoshimum, Y. Fujita, H. Toyokawa, K. Hosono and M. Kawabata	
Closing Session	
Concluding remarks	437
C. Glashausser	
Poster session	
The ($t, ^3\text{He}$) measurement at intermediate energy at NSCL	453
I. Daito, H. Akimune, S. M. Austin, D. Bazin, C. P. A. Berg, J. A. Brown, B. S. Davids, Y. Fujita, H. Fujimura, M. Fujiwam, R. Hazama, T. Inomata, K. Ishibashi, J. Janecke, S. Nakayama, K. Pham, D. A. Roberts, B. M. Sherrill, M. Steiner, A. Tamii, M. Tanaka, H. Toyokawa and M. Yosoi	
Microscopic structure of spin-isospin excited states in ^{13}N	456
H. Fujimura, H. Akimune, A. M. van den Berg, I. Daito, Y. Fujita, M. Fujiwara, M. N. Hamkeh, F. Ihara, T. Inomata, K. Ishibashi, J. Janecke, H. Kohri, H. K. T. van der Molen, S. Nakayama, A. Tamii, M. Tanaka, H. Toyokawa and M. Yosoi	
Search for annual modulation of WIMPs by means of large volume NaI(Tl) detector in ELEGANT V	459
K. Fushimi, H. Ejiri, K. Hayashi, N. Kudomi, K. Kume, K. Takahisa, S. Yoshida, R. Hazama, T. Kishimoto, M. Komori, K. Matsuoka, H. Miyawaki, S. Shiomi and H. Ohsumi	
Study of ^{28}Si , ^{32}S and $^{34}\text{S}(d, ^2\text{He})$ at $E_d = 270$ MeV	462
K. Itoh, T. Niizeki, T. Ichiham, H. Sakai, H. Okamura, S. Ishida, N. Sakamoto, T. Uesaka, Y. Satou, T. Ohnishi, Y. Tajima, G. Yokoyama, K. Sekiguchi, K. Yako, S. Fukusaka, H. Ohnuma and H. Oriham	

Measurements of the polarization transfer observables for the zero-degree $^{28}\text{Si}(\mathbf{p}, \mathbf{p}')$ reactions	465
<u>T. Kawabata, T. Ishikawa, M. Itoh, M. Nakamura, H. Sakaguchi,</u>	
<u>T. Taki, H. Takeda, A. Tamii, M. Yosoi, H. Akimune, I. Daito</u>	
<u>M. Fujiwara, K. Hatanaka, F. Ihara, T. Inomata, K. Ishibashi,</u>	
<u>T. Noro, E. Obayashi, H. Yoshida, M. Yoshimum, Y. Fujita,</u>	
<u>K. Hosono, H. Toyokawa and M. Kawabata</u>	
Study of reaction mechanisms for (\mathbf{p}, \mathbf{n}) and $(\mathbf{p}, \mathbf{p}')$ quasi elastic scattering	468
<u>H. Otsu, H. Sakai, H. Okamura, T. Wakasa, K. Hatanaka,</u>	
<u>S. Ishida, N. Sakamoto, T. Uesaka, Y. Satou, S. Fujita,</u>	
<u>A. Okihana, N. Koori and M. B. Greenfield</u>	
Total widths of isobaric analog states in ^{140}Pr , $^{172,174,176}\text{Lu}$ and ^{208}Bi	471
<u>A. Terakawa, C. C. Yun, H. Orihara, Y. Tajima, K. S. Itoh,</u>	
<u>A. Yamamoto, N. Matsumura, K. Kawami, H. Suzuki,</u>	
<u>G. Kamurai, H. Mizuno, Y. Fujii, T. Aizawa, M. Matsunaga,</u>	
<u>M. Hirai, T. Nakagawa and K. Ishii</u>	
Spin-isospin excitations via the $(^6\text{Li}, ^6\text{He})$ reaction	474
<u>H. Ueno, Y. Fujita, H. Fujita, Y. Kanzaki, T. Shinada,</u>	
<u>K. Katori, M. Yosoi, A. Tamii, F. Ihara, K. Takahisa,</u>	
<u>H. Kohri and T. Yamagata</u>	
Study of deuteron high momentum component via the ${}^3\text{He}(\vec{d}, p){}^4\text{He}$ reaction at 270MeV	477
<u>T. Uesaka, H. Sakai, H. Okamura, T. Ohnishi, Y. Satou, H. Otsu,</u>	
<u>T. Wakasa, T. Nonaka, G. Yokoyama, K. Sekiguchi, S. Haruyama,</u>	
<u>T. Ikeda, S. Ishida, N. Sakamoto, Y. Yano, T. Wakui and K. Itoh</u>	
Study of effective NN interaction in nuclei via the $^{28}\text{Si}(\vec{p}, \vec{n}){}^{28}\text{P}(6^-)$ reaction at 295MeV	480
<u>K. Yako, H. Sakai, H. Okamura, T. Nonaka, S. Fukusaka,</u>	
<u>K. Sekiguchi, T. Wakasa, H. Otsu and K. Hatanaka</u>	
Study of isoscalar spin excitations via the $(\mathbf{d}, \mathbf{d}')$ reaction	483
<u>Y. Satou, H. Sakai, H. Okamura, T. Nonaka, T. Ohnishi,</u>	
<u>G. Yokoyama, K. Sekiguchi, K. Yako, S. Fukusaka, S. Ishida,</u>	
<u>H. Otsu, N. Sakamoto, T. Uesaka, T. Wakasa, T. Ichihara,</u>	
<u>K. Itoh, T. Niizeki and K. Hatanaka</u>	

An experimental ensemble for the polarization transfer measurements in the (p, p') scattering at zero degrees	486
<i>M. Yosoi, H. Akimune, I. Daito, Y. Fujita, M. Fujiwara, K. Hatanaka, K. Hosono, F. Ihara, T. Inomata, K. Ishibashi, T. Ishikawa, M. Itoh, M. Kawabata, T. Kawabata, M. Nakamura, T. Noro, E. Obayashi, H. Sakaguchi, A. Tamii, H. Takeda, T. Taki, H. Toyokawa, H. Yoshida and M. Yoshimum</i>	
Spin-Isospin excitation observed in the (p, n) reaction on $^{6,7}\text{Li}$ at 35MeV	489
<i>H. Orihara, A. Temkawa, C. C. Yun, K. Itoh, A. Yamamoto, Y. Temmoto, N. Matsumura, G. C. Jon, T. Nakagawa, K. Ishii and H. Ohnuma</i>	
Program	497
List of Participants	503